

# REDEFINING SPINORS IN LORENTZ-VIOLATING QED

Don Colladay and Patrick McDonald

*New College of Florida*

*Sarasota, FL, 34243, U.S.A.*

An analysis of spinor redefinitions in the context of the Lorentz-violating QED extension is performed. Certain parameters that apparently violate Lorentz invariance are found to be physically irrelevant as they can be removed from the lagrangian using an appropriate redefinition of the spinor field components. It is shown that conserved currents may be defined using a modified action of the complex extension of the Lorentz group on the redefined spinors. This implies a natural correspondence between the apparently Lorentz-violating theory and conventional QED. Redefinitions involving derivatives are shown to relate certain terms in the QED extension to lagrangians involving nonlocal interactions or skewed coordinate systems. The redundant parameters in the QED extension are identified and the lagrangian is rewritten in terms of physically relevant coupling constants. The resulting lagrangian contains only physically relevant parameters and transforms conventionally under Lorentz transformations.

PACS: 11.30.Cp, 12.60.-i

## I. INTRODUCTION

The possibility of miniscule violations of Lorentz invariance arising from a more fundamental theory of nature has been of recent interest [1]. For example, such violations may arise in the low-energy limit of string theory [2], or physically realistic noncommutative field theories [3]. The full standard model extension uses the general concept of spontaneous symmetry breaking to construct a lagrangian consisting of all possible terms involving standard-model fields that are observer Lorentz scalars, including terms having coupling coefficients with Lorentz indices. At low energies, the relevant operators that are gauge invariant all have mass dimension  $D \leq 4$ , and are given in [4]. At higher energy scales nonrenormalizable terms are expected to play a role in the theoretical consistency of the model [5].

Various experiments have placed stringent bounds on parameters in the standard-model extension, including comparative tests of quantum electrodynamics (QED) in Penning traps and colliders [6, 7, 8, 9, 10, 11], spectroscopy of hydrogen and antihydrogen [12, 13], measurements of muon properties [14, 15], clock-comparison experiments [16, 17, 18, 19], observations of the behavior of a spin-polarized torsion pendulum [20, 21], measurements of cosmological birefringence [22, 4, 23, 24, 25], studies of neutral-meson oscillations [26, 27, 28], and observations of the baryon asymmetry [29].

In the theoretical results involving experimentally observable quantities, some of the parameters in the standard model extension do not appear while others occur only in specific linear combinations. The reason behind this is that some parameters that apparently violate Lorentz invariance when the spinor field is assumed to transform in the standard way under the action of the Lorentz group do not in fact violate this symmetry when the action on the field is appropriately modified such that the associated Lorentz currents are conserved. The freedom to select spinor coordinates in different ways generates a natural equivalence relation on the collection of lagrangians. Different lagrangians in the same equivalence class are related by field redefinitions; that is, by an invertible map between fields used to describe the same physics. The explicit construction of the redefinition used in this paper appears in Eq.(5). All

lagrangians in the same class are physically equivalent and the spinor transformation properties can be implemented so that the Lorentz currents are as close to conserved as possible. In other words, redefinitions may be used to define the currents so that they absorb the apparently Lorentz-violating terms which are obstructions to conservation. This means that one can use the transformation properties to eliminate a subset of the parameters appearing in the standard model extension and no more.

In this paper, the effects of field redefinitions in the context of extended QED are examined in detail. Particular terms in the standard model extension are already known to be unobservable since explicit redefinitions of the spinor components have previously been considered [4]. It is the goal of this work to analyze a more general set of field redefinitions and use them to simplify the full Lorentz-violating lagrangian as much as possible. The basic idea is to remove parameters that depend explicitly on the spinor coordinates. Once these redundant parameters have been eliminated, the remaining field transforms according to the standard action of the complex Lorentz group.

The paper is organized as follows. In section II, the extended QED theory is summarized. In Section III, an analysis is presented of the field redefinitions that are used to generate specific terms in the QED extension. The effects of transformations which do not include terms involving differentiation as well as those that do are investigated. The currents associated with U(1) and Poincaré group transformations for the general QED extension are derived in section IV. It is shown that conserved currents can be defined when only redundant parameters are present by using a similar representation to the conventional complex Lorentz group action. Section V contains the construction of the physical extended QED lagrangian with all redundant parameters removed.

## II. EXTENDED QED

To study the effects of specific field redefinitions, we restrict our attention to the QED subset involving only the electron and photon sectors of the full standard model extension presented in reference [4]. In the pure-photon sector, there is one CPT-even

$(k_F)$  and one CPT-odd ( $k_{AF}$ ) Lorentz-violating term. The free photon lagrangian is

$$\mathcal{L}_{\text{photon}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{4}(k_F)_{\kappa\lambda\mu\nu}F^{\kappa\lambda}F^{\mu\nu} + \frac{1}{2}(k_{AF})^\kappa\epsilon_{\kappa\lambda\mu\nu}A^\lambda F^{\mu\nu} \quad , \quad (1)$$

where the coupling  $k_F$  is a real, dimensionless coupling that can be taken to have the symmetries of the Riemann tensor, and the coefficient  $k_{AF}$  is real and has dimensions of mass.

Denoting the four-component electron field by  $\psi$  and the electron mass by  $m$ , the general QED lagrangian for electrons and photons including Lorentz-violating interactions arising from a generic spontaneous symmetry breaking mechanism is

$$\mathcal{L}_{\text{electron}}^{\text{QED}} = \frac{i}{2}\bar{\psi}\Gamma_\nu \overleftrightarrow{D}^\nu \psi - \bar{\psi}M\psi \quad , \quad (2)$$

where  $D_\mu = \partial_\mu + iqA_\mu$  is the usual covariant derivative,

$$\Gamma_\nu = \gamma_\nu + c_{\mu\nu}\gamma^\mu + d_{\mu\nu}\gamma_5\gamma^\mu + e_\nu + if_\nu\gamma_5 + \frac{1}{2}g_{\lambda\mu\nu}\sigma^{\lambda\mu} \quad , \quad (3)$$

and

$$M = m + im_5\gamma_5 + a_\mu\gamma^\mu + b_\mu\gamma_5\gamma^\mu + \frac{1}{2}H_{\mu\nu}\sigma^{\mu\nu} \quad . \quad (4)$$

Note that any Lorentz-*preserving* terms that arise from spontaneous symmetry breaking can be absorbed into the bare mass terms  $m$ ,  $m_5$ , and the overall normalization of the lagrangian. The normalization is chosen such that the coefficient of the  $\gamma_\nu$  term in Eq.(3) is one. The coupling coefficients  $a$ ,  $b$ ,  $c$ ,  $d$ ,  $e$ ,  $f$ ,  $g$ ,  $m_5$ , and  $H$  are real, constant parameters related to the vacuum expectation value of contributing tensor fields in the underlying theory.

Some of these parameters (or combinations of parameters) are only apparently Lorentz-violating as the lagrangian containing them can be shown to be equivalent to the standard lagrangian using the appropriate field redefinition. This question is taken up in the next section.

### III. GENERAL FERMION FIELD REDEFINITIONS

Some of the Lorentz-violating couplings in Eq.(2) can in fact be eliminated through a redefinition of the spinor field. To determine precisely which terms can be removed

in this manner, it is useful to begin with the standard Dirac lagrangian in terms of  $\psi$  with no Lorentz-violating terms and perform a field redefinition of the form  $\psi = R\chi$ , where  $R$  is some operator. The lagrangian in terms of  $\chi$  will contain terms included in the full Lorentz-violating lagrangian. In this section, we examine possible choices for the field redefinition and examine the resulting terms in the lagrangian.

To see which terms can be removed from the theory, we consider generic redefinitions of the fermion fields of the form

$$\psi(x) = [1 + f(x, \partial)]\chi(x) \quad , \quad (5)$$

where  $f(x, \partial)$  represents a general  $4 \times 4$  matrix function of the coordinates and derivatives. Only lowest order terms in the field redefinition parameters are retained since the Lorentz-violating couplings in the full lagrangian are assumed small. By applying this transformation to the conventional free fermion lagrangian (containing no Lorentz-violating parameters) we will see which terms can be eliminated from the extended theory by applying the inverse transformation.

To simplify the task, note that the Lorentz-violating terms generated by this transformation must have no explicit dependence on the coordinates and must consist of dimension  $D \leq 4$  operators<sup>1</sup>. Candidate terms for  $f(x, \partial)$  up to second order in  $x$  and  $\partial$  are of the form

$$f(x, \partial) = v \cdot \Gamma + i\theta + iA_\mu x^\mu + B_\mu \partial^\mu + \gamma_5 \tilde{B}_\mu \partial^\mu + C_{\mu\nu} x^\mu \partial^\nu \quad , \quad (6)$$

where  $v$  represents a set of arbitrary complex constants multiplying an arbitrary gamma matrix, denoted  $\Gamma$ , in the set  $m\{i\gamma_5, \gamma^\mu, \gamma_5\gamma^\mu, \sigma^{\mu\nu}\}$ ,  $\theta$  is a complex constant, while  $A_\mu$ ,  $B_\mu$ ,  $\tilde{B}_\mu$ , and  $C_{\mu\nu}$  are arbitrary real constants. Note that this is a generalization of the field redefinitions previously considered in [4].

The terms  $\text{Re}\theta$ ,  $B_\mu$  and the antisymmetric part of  $C_{\mu\nu}$  (together with the appropriate spin components  $v_{\mu\nu}$ ) are simply the generators of the  $U(1)$  and Poincaré groups and are symmetries of the conventional lagrangian. These terms do not generate any

---

<sup>1</sup>Actually, terms in the transformed lagrangian may contain an explicit dependence on  $x$  or may be of dimension greater than four provided that the terms are total divergences and therefore can be removed from the action. For examples, see the relevant terms in Eq.(11).

artificial Lorentz-violating parameters. The term  $\text{Im}\theta$  rescales the lagrangian and can be absorbed into the other constants. This leaves several independent transformations that may generate artificial Lorentz-violating terms. We proceed to calculate these explicitly in the rest of this section. Since we are working to lowest order in Lorentz-violating parameters, we can consider each term independently.

First, we summarize the results obtained using the  $v$  terms which have been previously described in [4]. An explicit example is presented to illustrate the general method. As is shown in the next section, a field redefinition of this type can be interpreted as selecting a new basis in spinor space for the representations of  $\text{SL}(2, \mathbb{C})$ , the complex extension of the Lorentz group. The standard lagrangian expressed in terms of the redefined field is given by

$$\begin{aligned}\mathcal{L} &= \frac{i}{2}\bar{\psi}\gamma^\mu \overleftrightarrow{\partial}_\mu \psi - m\bar{\psi}\psi \\ &= \mathcal{L}_0 + \frac{i}{2}\bar{\chi}[\{\gamma^\mu, \Gamma \cdot \text{Rev}\} + i[\gamma^\mu, \Gamma \cdot \text{Im}v]] \overleftrightarrow{\partial}_\mu \chi - 2m\text{Rev} \cdot \bar{\chi}\Gamma\chi \quad ,\end{aligned}\quad (7)$$

where  $\mathcal{L}_0$  is the conventional free field lagrangian in terms of  $\chi$ . For example, consider the field redefinition induced by  $v \cdot \Gamma = v_\mu \gamma^\mu$ . Using the above relation yields

$$\mathcal{L} = \mathcal{L}_0 + \text{Rev}_\mu [i\bar{\chi} \overleftrightarrow{\partial}^\mu \chi - 2m\bar{\chi}\gamma^\mu \chi] - i\text{Im}v_\mu [\bar{\chi}\sigma^{\mu\nu} \overleftrightarrow{\partial}_\nu \chi]. \quad (8)$$

Inspection of the term proportional to  $\text{Im}v_\mu$  indicates that the four terms in the extended lagrangian (2) of the form

$$g_{\lambda\mu\nu} = 2\text{Im}(v_\mu g_{\lambda\nu} - v_\lambda g_{\mu\nu}) \quad , \quad (9)$$

do not contribute in lowest order to the free fermion lagrangian. Examination of the terms multiplying  $\text{Rev}_\mu$  indicates that the simultaneous choice of

$$e^\mu = 2\text{Rev}^\mu \quad , \quad a^\mu = 2m\text{Rev}^\mu \quad (10)$$

can be removed from the lagrangian. This means that the field redefinition can remove either  $e^\mu$  or  $a^\mu$ , but not both, unless  $a^\mu = me^\mu$  happens to hold in the original lagrangian. Similar calculations can be done for the other choices of  $v \cdot \Gamma$ . The results are summarized in table 1. Note that a (finite) transformation of the form

| $\mathbf{v} \cdot \Gamma$   | $\mathbf{v} \in \Re \quad (v \equiv \text{Re } \mathbf{v})$                                                   | $\mathbf{v} \in \Im \quad (v \equiv \text{Im } \mathbf{v})$          |
|-----------------------------|---------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------|
| $v(i\gamma_5)$              | Used to eliminate $m_5$ term                                                                                  | $d_{\mu\nu} = 2vg_{\mu\nu}$                                          |
| $v_\mu\gamma^\mu$           | $e_\mu = 2v_\mu$ and $a_\mu = 2mv_\mu$                                                                        | $g_{\lambda\mu\nu} = 2(v_\mu g_{\lambda\nu} - v_\lambda g_{\mu\nu})$ |
| $v_\mu\gamma_5\gamma^\mu$   | $g_{\lambda\mu\nu} = -2\epsilon_{\lambda\mu\nu}{}^\alpha v_\alpha$ and $b_\mu = 2mv_\mu$                      | $f_\mu = -2v_\mu$                                                    |
| $v_{\mu\nu}\sigma^{\mu\nu}$ | $d_{\mu\nu} = -2\epsilon_{\mu\nu}{}^{\alpha\beta} v_{\alpha\beta}$ and $\frac{1}{2}H_{\mu\nu} = 2mv_{\mu\nu}$ | $c_{\mu\nu} = 2v_{[\mu\nu]}$                                         |

Table 1: Summary of terms generated by field redefinitions of the form  $v \cdot \Gamma$ .

$e^{iv\gamma_5}$  with  $v \in \Re$  is used to remove any term of the form  $m_5$  in the original lagrangian to all orders. The effect is an  $m_5$  dependent mixing of some of the Lorentz-violating parameters, but the structure is essentially unchanged.

Next, we consider the  $A$ ,  $B$ ,  $\tilde{B}$ , and  $C$  redefinitions. To lowest order in these parameters, the transformed lagrangian becomes

$$\begin{aligned} \mathcal{L} = \mathcal{L}_0 - \bar{\chi} A_\mu \gamma^\mu \chi + B_\mu \partial^\mu \mathcal{L}_0 - \tilde{B}_\mu \partial^\mu [\frac{i}{2} \bar{\chi} \gamma_5 \gamma^\nu \overleftrightarrow{\partial}_\nu \chi] - m \bar{\chi} \gamma_5 \tilde{B}_\mu \overleftrightarrow{\partial}^\mu \chi \\ + C_{\mu\nu} x^\mu \partial^\nu \mathcal{L}_0 + \frac{i}{2} C_{\mu\nu} \bar{\chi} \gamma^\mu \overleftrightarrow{\partial}^\nu \chi \quad . \end{aligned} \quad (11)$$

The  $A$  term can be used to eliminate  $a^\mu$  as is discussed in detail in reference [4]. The  $B$  term is a total divergence that drops out of the action. This is a direct consequence of translational invariance since under a (finite) translation of the coordinates by  $B$

$$\psi(x) = e^{B \cdot \partial} \chi(x) = \chi(x + B) = \chi(x') \quad , \quad (12)$$

and the action takes the same form in the translated coordinate system. The first  $C$  term can be partially integrated to yield a total divergence and a rescaling of  $\mathcal{L}_0$ . The final  $C$  term is of the form  $c$  as defined in Eq.(2).

A few remarks are in order regarding the above transformation involving  $C$ . First, note that such a field redefinition appears equivalent to changing fermion coordinates to a system with a new (constant) metric that skews the coordinates. We can see this by examination of the transformation

$$\psi(x) = (1 + C^{\mu\nu} x_\mu \partial_\nu) \chi(x) \approx e^{C^{\mu\nu} x_\mu \partial_\nu} \chi(x) = \chi(x + C \cdot x) = \chi(x') \quad , \quad (13)$$

where  $x'^\mu = x^\mu + C^\mu{}_\nu x^\nu$  are the new field coordinates. This redefinition is therefore equivalent to transforming to a skewed coordinate system with a nondiagonal metric given by

$$g'^{\mu\nu} = \eta^{\mu\nu} + C^{(\mu\nu)} \quad . \quad (14)$$

Rewriting the transformed lagrangian in terms of this new metric yields

$$\mathcal{L} = \frac{i}{2}\bar{\chi}(x')\tilde{\gamma}^\mu\overset{\leftrightarrow}{\partial}'_\mu\chi(x') - m\bar{\chi}(x')\chi(x') \quad (15)$$

where the modified matrices  $\tilde{\gamma}^\mu = (\eta^\mu_\nu + c^\mu_\nu)\gamma^\nu$  satisfy the relations  $\{\tilde{\gamma}^\mu, \tilde{\gamma}^\nu\} = 2g'^{\mu\nu}$ . The resulting lagrangian can be related to the conventional one using the vierbein formalism of general relativity by performing the appropriate general coordinate transformation. This shows that there is a natural association between the theory containing a  $c$  term and a theory formulated in a skewed coordinate system defined by the metric given above. If the free fermions are the only component to the theory it is possible to perform the appropriate general coordinate transformation on the skewed coordinates relating it to the conventional case. This is because it is not possible to distinguish the theory in a skewed coordinate system (with a  $c$  term) from a conventional theory in an orthonormal system since fermion propagation properties are the only tool available to define the coordinate system itself.

However, when photon interactions are incorporated through the covariant derivative, it is no longer possible to perform the general coordinate transformation without affecting the photon sector. There is now an alternate way to distinguish the coordinates physically so that the skewed coordinates become observable. Under the fermion field redefinition, the fermion-photon interaction term becomes

$$\mathcal{L}_{int} = -q\bar{\psi}(x)\gamma^\mu A_\mu(x)\psi(x) \longrightarrow -q\bar{\chi}(x')\tilde{\gamma}^\mu A'_\mu(x)\chi(x') \quad (16)$$

where the photon field is expressed as a function of the conventional coordinates  $x$ , but its components are resolved in the modified coordinates  $x'$ . In this picture, the theory exhibits a form of nonlocality since the fermion fields interact with the photon field at different spacetime points. If the photon field is re-expressed in terms of the new coordinates  $x'$ , the lagrangian becomes local, but picks up an extra term that breaks the natural association between the theories. Yet another approach is to redefine the physical photon field  $A'_\mu(x) \rightarrow A'_\mu(x')$ , but the new metric introduces corrections of the form in Eq.(1) into the kinetic photon sector. Therefore, the photon interactions prevent the trivial elimination of symmetric  $c$  terms using the above field redefinition. Similar problems arise when using other derivative transformations, so these are not

considered in detail in subsequent sections of this paper. For example, the term  $\tilde{B}$  in Eq.(6) generates a term of the form  $f$  (defined in Eq.(2)) as well as a total divergence in the transformed lagrangian of Eq.(11). This transformation corresponds to a shift in opposite directions for the left-handed and right-handed fields due to the presence of  $\gamma_5$ . There is therefore a natural correspondence between the lagrangian with an  $f$  term and the conventional theory provided the left-handed and right-handed fields can be translated independently<sup>2</sup>. The interaction that breaks the correspondence with the conventional theory in this case is the fermion mass term that mixes left-handed and right-handed fields. A similar situation occurs when the  $C$  transformation discussed above is multiplied by  $\gamma_5$ . Symmetric components of the form  $d$  defined in Eq.(2) arise in the transformed lagrangian, but the mass term depends explicitly on  $x$ , therefore breaking the natural correspondence between the theories.

Finite field transformations can be constructed through exponentiation of Eq.(5). The results of these transformations are often much more complicated than the infinitesimal ones since several parameters can be mixed. As an example, consider a field redefinition of the form

$$\psi = e^{v_\mu \gamma^\mu} \chi = (\cosh v + \frac{v_\mu \gamma^\mu}{v} \sinh v) \chi \quad , \quad (17)$$

with  $v_\mu$  real, timelike, and the quantity  $v$  defined by  $v = \sqrt{v_\mu v^\mu}$ . Application of this transformation to the standard lagrangian in terms of  $\psi$  yields a lagrangian for  $\chi$  with apparent Lorentz-violating parameters and a modified mass given by

$$\begin{aligned} e^\mu &= \frac{v^\mu}{v} \sinh 2v \quad ; \quad c^{\mu\nu} = \frac{v^\mu v^\nu}{v^2} (\cosh 2v - 1) \quad , \\ m' &= m \cosh 2v \quad ; \quad a^\mu = \frac{m v^\mu}{v} \sinh 2v \quad . \end{aligned} \quad (18)$$

Note that the corrections to  $c$  and  $m$  terms appear only at second order in  $v$ .

Such a choice of parameters in the QED extension leaves the dispersion relation unaltered and therefore leads to no stability problems or microcausality violations. This is true for any finite field redefinition of the form  $e^{v \cdot \Gamma}$  since the field redefinition

---

<sup>2</sup>Actually, one of the redefinitions listed in table 1 can also be used to eliminate the  $f$  term, so it is always removable from the lagrangian.

commutes with the square of the conventional Dirac equation. The set of all such transformations generates a class of lagrangians equivalent to the conventional one.

The derivative field redefinitions may also be exponentiated to yield finite transformations. In the cases of  $\tilde{B}$  and  $C$  of Eq.(6), finite transformations correspond to finite coordinate transformations, possibly different for various spinor components. As in the infinitesimal case, interactions between various spinor components and other fields limit the usefulness of these transformations due to nonlocality problems.

To gain further insight into the invariance of the physics under the above transformations, it is useful to compute the currents associated with the generators of the Poincaré group. We will see that it is necessary to redefine the action of the complex Lorentz group along with the field in order to yield maximally conserved Lorentz generators.

#### IV. POINCARÉ GENERATORS

We start with a general free fermion lagrangian of the form

$$\mathcal{L} = \frac{i}{2} \bar{\chi} \Gamma^\nu \overleftrightarrow{\partial}_\nu \chi - \bar{\chi} M \chi \quad , \quad (19)$$

and apply Noether's theorem to obtain the divergence of the currents associated with various continuous transformations of the field. Invariance under a global  $U(1)$  phase transformation yields a conserved current of

$$j^\mu = \bar{\chi} \Gamma^\mu \chi \quad , \quad (20)$$

satisfying  $\partial_\mu j^\mu = 0$ . Invariance under translations yields a conserved energy momentum tensor of

$$\Theta^{\mu\nu} = \frac{i}{2} \bar{\chi} \Gamma^\mu \overleftrightarrow{\partial}^\nu \chi \quad , \quad (21)$$

satisfying  $\partial_\mu \Theta^{\mu\nu} = 0$ .

The lagrangian is no longer invariant under the conventional action of the Lorentz group, so the divergences of the corresponding currents will not vanish. These can be calculated using the standard technique of writing the action over an arbitrary four-volume in terms of boosted coordinates and fields at  $x'^\mu = \Lambda^\mu_\nu x^\nu \approx x^\mu + \epsilon^\mu_\nu x^\nu$

and calculating the variation to lowest order in  $\epsilon^\mu_{\nu}$ . A choice must be made for the induced transformation properties of the spinor components of  $\chi$  under the complex extension of the Lorentz group  $SL(2, C)$ . Using the standard  $S(\Lambda) = 1 - \frac{i}{4}\sigma_{\mu\nu}\epsilon^{\mu\nu}$  yields currents given by

$$\partial_\alpha j^{\alpha\mu\nu} = X^{\mu\nu} \quad , \quad (22)$$

where

$$j^{\alpha\mu\nu} = x^{[\mu}\Theta^{\alpha\nu]} + \frac{1}{4}\bar{\chi}\{\Gamma^\alpha, \sigma^{\mu\nu}\}\chi \quad , \quad (23)$$

and

$$\begin{aligned} X^{\mu\nu} = & -a^{[\mu}\bar{\chi}\gamma^{\nu]}\chi - b^{[\mu}\bar{\chi}\gamma_5\gamma^{\nu]}\chi - \frac{i}{2}\bar{\chi}(c^{[\nu\alpha}\gamma^{\mu]}\overset{\leftrightarrow}{\partial}_\alpha + c^{\alpha[\nu}\gamma_\alpha\overset{\leftrightarrow}{\partial}^{\mu]})\chi \\ & - \frac{i}{2}\bar{\chi}(d^{[\nu\alpha}\gamma_5\gamma^{\mu]}\overset{\leftrightarrow}{\partial}_\alpha + d^{\alpha[\nu}\gamma_5\gamma_\alpha\overset{\leftrightarrow}{\partial}^{\mu]})\chi + \frac{i}{2}e^{[\mu}\bar{\chi}\overset{\leftrightarrow}{\partial}^{\nu]}\chi \\ & - \frac{1}{2}f^{[\mu}\bar{\chi}\gamma_5\overset{\leftrightarrow}{\partial}^{\nu]}\chi - \frac{i}{4}\bar{\chi}(2g^{[\nu\alpha\beta}\sigma^\mu_\alpha\overset{\leftrightarrow}{\partial}_\beta + g^{\alpha\beta[\nu}\sigma_{\alpha\beta}\overset{\leftrightarrow}{\partial}^{\mu]})\chi - \bar{\chi}H^{[\mu\alpha}\sigma^\nu_{\alpha}\chi \quad . \quad (24) \end{aligned}$$

All of these equations can be verified by direct calculation.

It is expected that terms which can be eliminated using a field redefinition should also be removable from  $X^{\mu\nu}$  with the appropriate redefinition of the associated currents. To explicitly construct the conserved currents, the field redefinition is applied to the conventional currents written in terms of  $\psi$ , satisfying the conventional Dirac equation. These conserved currents are re-expressed in terms of  $\chi$  giving the proper conserved currents for the new lagrangian.

Alternatively, the conserved currents of the modified lagrangian may be computed from Nöether's theorem using a modified action of the Lorentz group on the spinor fields. Using a general finite field redefinition  $\psi = e^{v\cdot\Gamma}\chi$ , it is found that the action of U(1) and translations is the same on  $\psi$  and  $\chi$  meaning that the 4-current and energy-momentum tensors are computed as in Eqs.(20, 21). However, the action of the Lorentz transformations are in fact modified due to the field redefinition. Under an infinitesimal Lorentz transformation of the coordinates, the  $\psi$  spinor components mix according to  $\delta\psi = S(\Lambda)\psi$ , while the corresponding change in  $\chi$  is calculated to be  $\delta\chi = e^{-v\cdot\Gamma}S(\Lambda)e^{v\cdot\Gamma}\chi = \tilde{S}(\Lambda)\chi$ . This means that the  $\chi$  components transform according to a similar representation of the complex Lorentz group. When the asso-

ciated current is computed using this modified action of  $SL(2, \mathbb{C})$  on  $\chi$ , it is indeed conserved.

As an example, consider the field redefinition used in Eq.(8) involving  $Rev_\mu$ . If the conventional  $S(\Lambda)$  is used to find the current associated with the Lorentz transformations, the result is  $j_\chi^{\alpha\mu\nu}$  given by Eq.(23) which is not conserved. However, if  $\tilde{S}(\Lambda)$  is used to transform the field  $\chi$ , then the associated current is

$$\tilde{j}_\chi^{\alpha\mu\nu} = x^{[\mu}\Theta^{\alpha\nu]} + \frac{1}{4}\bar{\chi}\{\Gamma^\alpha, \sigma^{\mu\nu}\}\chi + \frac{1}{2}\bar{\chi}e^{[\mu}\sigma^{\nu]\alpha}\chi \quad , \quad (25)$$

which is in fact conserved. This shows that one must be careful to map the correct conserved generators into the proper associated currents written in terms of  $\chi$ . Similar maps between generators can be performed using the derivative transformations, however, in this case the  $U(1)$  and translation currents may also be modified.

As a practical approach, the redundant parameters can first be removed from the lagrangian and then the conventional  $S(\Lambda)$  can be used to construct the currents in the redefined lagrangian. We perform this removal of the redundant parameters in the next section.

## V. ELIMINATION OF REDUNDANT PARAMETERS

In this section, we start with a general lagrangian and apply field redefinitions to remove redundant parameters. The starting point is the lagrangian given in Eq.(2). The parameters  $m_5$ ,  $a_\mu$ ,  $e_\mu$ ,  $f_\mu$ , and  $c_{[\mu\nu]}$  can all be immediately removed using a combination of the transformations described in section III. The parameter  $g_{\lambda\mu\nu}$  can be replaced by the traceless  $\tilde{g}_{\lambda\mu\nu}$  satisfying  $g^{\lambda\nu}\tilde{g}_{\lambda\mu\nu} = g^{\mu\nu}\tilde{g}_{\lambda\mu\nu} = 0$ . The other transformations listed in table 1 involve linear combinations of parameters.

To handle these terms, the lagrangian must be re-expressed in terms of the new, physically relevant linear combinations of parameters. For example, the combination of antisymmetric  $d$  and  $H$  terms of Eq.(2) are re-expressed as

$$\begin{aligned} \frac{i}{4}\bar{\psi}d_{[\mu\nu]}\gamma_5\gamma^\mu\overleftrightarrow{\partial}^\nu\psi - \frac{1}{2}\bar{\psi}H_{\mu\nu}\sigma^{\mu\nu}\psi = & v_{\alpha\beta}^+ \left[ \frac{i}{2}\bar{\psi}\epsilon_{\mu\nu}^{\alpha\beta}\gamma_5\gamma^\mu\overleftrightarrow{\partial}^\nu\psi - m\bar{\psi}\sigma^{\alpha\beta}\psi \right] \\ & + v_{\alpha\beta}^- \left[ \frac{i}{2}\bar{\psi}\epsilon_{\mu\nu}^{\alpha\beta}\gamma_5\gamma^\mu\overleftrightarrow{\partial}^\nu\psi + m\bar{\psi}\sigma^{\alpha\beta}\psi \right] \quad , \end{aligned} \quad (26)$$

where  $v_{\alpha\beta}^{\pm} = \frac{1}{4}(\tilde{d}_{\alpha\beta} \pm \frac{1}{m}H_{\alpha\beta})$ , and  $\epsilon_{\mu\nu}^{\alpha\beta}\tilde{d}_{\alpha\beta} = d_{[\mu\nu]}$ . The combination  $v_{\alpha\beta}^{-}$  can be removed using a field redefinition, leaving only  $v_{\alpha\beta}^{+}$  terms in the lagrangian. The  $g$  and  $b$  terms can be similarly combined using the definition  $v_{\alpha}^{\pm} = \frac{1}{2}(g_{\alpha} \pm \frac{1}{m}b_{\alpha})$

After all of these redefinitions have been performed, the form for the lagrangian becomes

$$\begin{aligned} \mathcal{L} = & \frac{i}{2}\bar{\psi}(\gamma_{\nu} + \frac{1}{2}c_{(\mu\nu)}\gamma^{\mu} + \frac{1}{2}d_{(\mu\nu)}\gamma_5\gamma^{\mu} + \frac{1}{2}\hat{g}_{\lambda\mu\nu}\sigma^{\lambda\mu})\overleftrightarrow{\partial}^{\nu}\psi - m\bar{\psi}\psi \\ & + \frac{1}{4}(\tilde{d}_{\alpha\beta} + \frac{1}{m}H_{\alpha\beta})\left[\frac{i}{2}\bar{\psi}\epsilon_{\mu\nu}^{\alpha\beta}\gamma_5\gamma^{\mu}\overleftrightarrow{\partial}^{\nu}\psi - m\bar{\psi}\sigma^{\alpha\beta}\psi\right] \\ & - \frac{1}{2}(g_{\alpha} - \frac{1}{m}b_{\alpha})\left[\frac{i}{2}\bar{\psi}\frac{1}{2}\epsilon_{\lambda\mu}^{\nu\alpha}\sigma^{\lambda\mu}\overleftrightarrow{\partial}_{\nu}\psi - m\bar{\psi}\gamma_5\gamma^{\alpha}\psi\right] \quad , \end{aligned} \quad (27)$$

where  $\hat{g}_{\lambda\mu\nu}$  is a traceless coupling with a vanishing totally antisymmetric piece. The totally antisymmetric component of  $\tilde{g}$  is absorbed into  $g_{\alpha} = -\epsilon_{\alpha}^{\lambda\mu\nu}\tilde{g}_{\lambda\mu\nu}$ . This lagrangian can be written in the form

$$\mathcal{L} = \frac{i}{2}\bar{\psi}\tilde{\Gamma}_{\nu}\overleftrightarrow{\partial}^{\nu}\psi - \bar{\psi}\tilde{M}\psi \quad , \quad (28)$$

where

$$\tilde{\Gamma}_{\nu} = \gamma_{\nu} + \frac{1}{2}c_{(\mu\nu)}\gamma^{\mu} + (\frac{1}{2}d_{(\mu\nu)} + \epsilon_{\mu\nu}^{\alpha\beta}v_{\alpha\beta}^{+})\gamma_5\gamma^{\mu} + \frac{1}{2}(\hat{g}_{\lambda\mu\nu} - \epsilon_{\lambda\mu\nu}^{\alpha}v_{\alpha}^{-})\sigma^{\lambda\mu} \quad , \quad (29)$$

and

$$\tilde{M} = m(1 + v_{\alpha\beta}^{+}\sigma^{\alpha\beta} - v_{\alpha}^{-}\gamma_5\gamma^{\alpha}) \quad , \quad (30)$$

where all distinct parameters are now physically relevant.

This lagrangian is therefore the one for which  $\psi$  can be assumed to transform under the standard representation of  $SL(2,C)$ , yielding maximally conserved currents<sup>3</sup>. The relevant currents are given as in the previous section with appropriately mapped constants found by comparison of Eq.(2) and Eq.(28). The remaining terms cannot be removed by a redefinition of spinor coordinates. Other terms may be eliminated in the free theory using transformations involving derivatives of the fields as is discussed in section III, but interactions between the fermion fields and other fields often break

---

<sup>3</sup>Note that field redefinitions involving the physically relevant coupling constants may still be performed yielding different definitions of the Lorentz currents. However, the physically irrelevant parameters have been removed. For example, a further field redefinition is convenient in removing extra time derivatives for calculational purposes as is described preceding Eq.(32).

the natural correspondence between the redefined lagrangian and the original theory. This means that if these terms are removed from the free fermion sector, they will appear as modified interaction terms and will not be removed from the theory, only shifted to another sector.

Physical quantities should therefore depend only on the combinations occurring in the lagrangian of Eq.(28). Therefore, it is not possible to obtain experimental bounds on all Lorentz-violating parameters of Eq.(2) independently, only on the combinations present in Eq.(28). For example, it is only the linear combination  $v_{\alpha\beta}^+$  of the anti-symmetric part of  $d$  and  $H$  that is observable. This implies that only one parameter should in fact be used to describe this quantity.

Comparison with previous calculations within the context of this model verify that this is indeed the case. For example, in applications to electrons and positrons in Penning traps [6, 7, 8, 9], the relevant experimental bound is obtained from the observable cyclotron and anomaly frequencies ( $\tilde{g}_{\lambda\mu\nu} = 0$  in this calculation)

$$\begin{aligned}\omega_c^{e\pm} &\approx (1 - c_{00}^e - c_{11}^e - c_{22}^e)\omega_c \quad , \\ \omega_c^{e\pm} &\approx \omega_a \mp 2b_3^e + 2md_{30}^e + 2H_{12}^e \quad ,\end{aligned}\tag{31}$$

which can place bounds only on the combinations of parameters found in Eq.(27), not on the parameters that can be removed by the field redefinitions. As another example, a calculation of the cross-section for  $e^+e^- \rightarrow 2\gamma$  within the QED extension [10] only depends on the symmetric components of  $c$ . Similar results have been obtained in other QED systems [30].

For many practical calculations it is convenient to perform another field redefinition to normalize  $\tilde{\Gamma}^0 \rightarrow \gamma^0$ . This has the effect of removing extra time derivative couplings insuring that the resulting Schrödinger Equation has a conventional time evolution [6, 17]. Starting with a general lagrangian of the form (2), the appropriate field redefinition that removes time derivative couplings to lowest order is  $\psi = A\chi$  with

$$\begin{aligned}A &= 1 - \frac{1}{2}\gamma^0(\Gamma_0 - \gamma_0) \\ &= 1 - \frac{1}{2}\gamma^0(c_{\mu 0}\gamma^\mu + d_{\mu 0}\gamma_5\gamma^\mu + e_0 + if_0\gamma_5 + \frac{1}{2}g_{\lambda\mu 0}\sigma^{\lambda\mu}) \quad .\end{aligned}\tag{32}$$

Note that these are in 1-1 correspondence with the non-unitary transformations of the form  $v \cdot \Gamma$  examined in section III. The unitary transformations of this type can be obtained by letting the coupling constants in Eq.(32) be pure imaginary rather than real. In this case, the time derivative structure is unaffected by the field redefinition. This provides an alternative perspective on the spinor component field redefinitions. The unitary transformations preserve the time derivative terms while the non-unitary transformations modify the time derivative structure. This can be seen directly from Eq.(7) by noting that the requirement that no time derivatives are introduced is equivalent to the condition  $(v \cdot \Gamma)^\dagger = -v \cdot \Gamma$ , hence making the field redefinition unitary to lowest order. The unitary transformations are used in the standard Dirac theory to transition between various representations of the gamma matrices, an alternative perspective to the explicit correction terms used in this paper. In other words, any apparent Lorentz-violating terms generated by a unitary transformation may be absorbed into a modified gamma matrix representation.

As a practical method for calculation, the procedure is therefore to first remove the redundant parameters to obtain Eq.(28), then perform the field redefinition of Eq.(32) (using  $\tilde{\Gamma}^0$  in place of  $\Gamma^0$ ). The resulting lagrangian will therefore yield a conventional Schrödinger Equation time evolution and will not contain any redundant parameters.

## VI. SUMMARY

In this paper, an analysis of field redefinitions in the context of the Lorentz-violating QED extension was presented. It was shown that a variety of parameters that apparently violate Lorentz invariance can be eliminated from the lagrangian using suitable fermionic field redefinitions.

The approach taken to find these parameters was to begin with the conventional Dirac lagrangian, introduce an arbitrary spinor redefinition, and examine the resulting transformed lagrangian. Any parameters generated using this procedure can be removed by the corresponding inverse transformation. The action of  $SL(2, \mathbb{C})$  on the spinor fields is deduced using the conventional action on the  $\psi$  spinors and performing the field redefinition to determine the corresponding action on  $\chi$ . The resulting trans-

formation matrices of the new spinor are related by a similarity transformation to the original matrices. This modified action must be taken into account when defining the conserved generators in the modified lagrangian.

This implies a procedure for elimination of redundant parameters from the original lagrangian. By first identifying all possible terms that can be generated from the conventional lagrangian by a field redefinition, these terms can be omitted from the physical lagrangian. This procedure has been carried out in section V of the paper. A further transformation may be implemented to normalize the time derivative structure of the theory to obtain conventional Schrödinger time evolution and physical particle states.

The possible nonderivative field redefinitions that can be applied to the QED extension fall into two general categories, unitary and nonunitary. The nonunitary transformations modify the time derivative structure of the lagrangian and can be used to construct a hermitian hamiltonian and a Schrödinger equation with conventional time evolution. The time derivative couplings in a general bilinear lagrangian of the form in Eq.(2) can always be removed using a suitable nonunitary transformation. Moreover, there is a one-to-one correspondence between these nonunitary transformations and the field redefinitions used to eliminate the extra time derivative couplings.

Redefinitions involving differentiation may be useful in the free fermion theory, but often lead to nonlocal interactions or skewed coordinate systems when interactions are present. These problems make it difficult to perform a generic analysis of all possible applications of these transformations. Derivative redefinitions may be applied on a case by case basis where they might be useful in simplification of calculations.

Stability and causality issues [5] cannot be effectively addressed using the above arguments since the redefinition was only carried out to lowest order in Lorentz-violating parameters. Causality and stability problems appear either when the coupling constants are large, or when the momentum is significantly large to invalidate the linear approximations involved. However, the finite field redefinition considered in Eq.(17) leads to a finite set of parameters that maintain the conventional dispersion relation. The resulting theory must therefore be stable and microcausal. A class of apparently

Lorentz-violating lagrangians that are microcausal and stable may therefore be generated by applying finite versions of the field redefinitions discussed in this paper. A complete nonperturbative analysis would be of interest, but is beyond the scope of this work.

Application of a similar analysis to the entire standard model extension would be of interest. For example, cross-generational mixings would allow for a richer structure of possible field redefinitions than in QED.

**ACKNOWLEDGMENTS** This work was supported in part by a University of South Florida Division of Sponsored Research grant. We thank Alan Kostelecký and Robert Bluhm for discussions.

## REFERENCES

1. For a summary of recent theoretical models and experimental tests see, for example, *CPT and Lorentz Symmetry*, V.A. Kostelecký, ed., World Scientific, Singapore, 1999; *CPT and Lorentz Symmetry II*, V.A. Kostelecký, ed., World Scientific, Singapore, 2002.
2. V.A. Kostelecký and S. Samuel, Phys. Rev. D **39**, 683 (1989); *ibid.* **40**, 1886 (1989); Phys. Rev. Lett. **63**, 224 (1989); *ibid.* **66**, 1811 (1991); V.A. Kostelecký and R. Potting, Nucl. Phys. B **359**, 545 (1991); Phys. Lett. B **381**, 89 (1996); Phys. Rev. D **63**, 046007 (2001); V.A. Kostelecký, M. Perry, and R. Potting, Phys. Rev. Lett. **84**, 4541 (2000).
3. S. Carroll, *et al.*, Phys. Rev. Lett. **87**, 141601 (2001).
4. D. Colladay and V.A. Kostelecký, Phys. Rev. D **55**, 6760 (1997); Phys. Rev. D **58**, 116002 (1998).
5. V.A. Kostelecký and R. Lehnert, Phys. Rev. D **63**, 065008 (2001).
6. R. Bluhm *et al.*, Phys. Rev. Lett. **79**, 1432 (1997); Phys. Rev. D **57**, 3932 (1998).
7. G. Gabrielse *et al.*, in Ref. [1]; Phys. Rev. Lett. **82**, 3198 (1999).
8. H. Dehmelt *et al.*, Phys. Rev. Lett. **83**, 4694 (1999).

9. R. Mittleman, I. Ioannou, and H. Dehmelt, in Ref. [1]; R. Mittleman *et al.*, Phys. Rev. Lett. **83**, 2116 (1999).
10. D. Colladay, and V.A. Kostelecký, Phys. Lett. B **511**, 209 (2001).
11. V.A. Kostelecký, C. D. Lane, and A.G. Pickering, To appear in Phys. Rev. D, hep-th/0111123.
12. R. Bluhm *et al.*, Phys. Rev. Lett. **82**, 2254 (1999).
13. D.F. Phillips *et al.*, Phys. Rev. D **63**, 111101 (2001).
14. R. Bluhm *et al.*, Phys. Rev. Lett. **84**, 1098 (2000).
15. V.W. Hughes *et al.*, presented at the Hydrogen II Conference, Tuscany, Italy, June, 2000.
16. V.W. Hughes, H.G. Robinson, and V. Beltran-Lopez, Phys. Rev. Lett. **4**, 342 (1960); R.W.P. Drever, Philos. Mag. **6**, 683 (1961); J.D. Prestage *et al.*, Phys. Rev. Lett. **54**, 2387 (1985); S.K. Lamoreaux *et al.*, Phys. Rev. Lett. **57**, 3125 (1986); Phys. Rev. A **39**, 1082 (1989); T.E. Chupp *et al.*, Phys. Rev. Lett. **63**, 1541 (1989); C.J. Berglund *et al.*, Phys. Rev. Lett. **75**, 1879 (1995).
17. V.A. Kostelecký and C.D. Lane, Phys. Rev. D **60**, 116010 (1999); J. Math. Phys. **40**, 6245 (1999).
18. L.R. Hunter *et al.*, in Ref. [1].
19. D. Bear *et al.*, Phys. Rev. Lett. **85**, 5038 (2000).
20. R. Bluhm and V.A. Kostelecký, Phys. Rev. Lett. **84**, 1381 (2000).
21. B. Heckel *et al.*, in B.N. Kursunoglu, S.L. Mintz, and A. Perlmutter, eds., *Elementary Particles and Gravitation*, Plenum, New York, 1999.
22. S.M. Carroll, G.B. Field, and R. Jackiw, Phys. Rev. D **41**, 1231 (1990).
23. R. Jackiw and V.A. Kostelecký, Phys. Rev. Lett. **82**, 3572 (1999).
24. M. Pérez-Victoria, Phys. Rev. Lett. **83**, 2518 (1999); J.M. Chung, Phys. Lett. B **461**, 138 (1999).
25. V.A. Kostelecký and M. Mewes, Phys. Rev. Lett. **87**, 251304 (2001).
26. V.A. Kostelecký and R. Potting, Phys. Rev. D **51**, 3923 (1995); D. Colladay

- and V.A. Kostelecký, Phys. Lett. B **344**, 259 (1995); Phys. Rev. D **52**, 6224 (1995); V.A. Kostelecký and R. Van Kooten, Phys. Rev. D **54**, 5585 (1996); V.A. Kostelecký, Phys. Rev. Lett. **80**, 1818 (1998); Phys. Rev. D **61**, 016002 (2000); hep-ph/0104120; N. Isgur *et al.*, in preparation.
27. KTeV Collaboration, Y.B. Hsiung *et al.*, Nucl. Phys. Proc. Suppl. **86**, 312 (2000).
  28. OPAL Collaboration, R. Akerstaff *et al.*, Z. Phys. C **76**, 401 (1997); DELPHI Collaboration, M. Feindt *et al.*, preprint DELPHI 97-98 CONF 80 (July 1997); BELLE Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **86**, 3228 (2001).
  29. O. Bertolami *et al.*, Phys. Lett. B **395**, 178 (1997).
  30. For a recent review of Lorentz tests in QED, see R. Bluhm, proceedings QED 2000, *Fronteir Tests of Quantum Electrodynamics and Physics of the Vacuum*, Trieste, Italy, October, 2000. (hep/ph-0011272).